

The age incidence of chronic myeloid leukemia can be explained by a one-mutation model

Franziska Michor, Yoh Iwasa, and Martin A. Nowak

PNAS 2006;103:14931-14934; originally published online Sep 25, 2006;
doi:10.1073/pnas.0607006103

This information is current as of October 2006.

Online Information & Services	High-resolution figures, a citation map, links to PubMed and Google Scholar, etc., can be found at: www.pnas.org/cgi/content/full/103/40/14931
References	This article cites 27 articles, 9 of which you can access for free at: www.pnas.org/cgi/content/full/103/40/14931#BIBL This article has been cited by other articles: www.pnas.org/cgi/content/full/103/40/14931#otherarticles
E-mail Alerts	Receive free email alerts when new articles cite this article - sign up in the box at the top right corner of the article or click here .
Rights & Permissions	To reproduce this article in part (figures, tables) or in entirety, see: www.pnas.org/misc/rightperm.shtml
Reprints	To order reprints, see: www.pnas.org/misc/reprints.shtml

Notes:

The age incidence of chronic myeloid leukemia can be explained by a one-mutation model

Franziska Michor*[†], Yoh Iwasa*[‡], and Martin A. Nowak[§]

[§]Program for Evolutionary Dynamics, Departments of Mathematics and Organismic and Evolutionary Biology, and *Society of Fellows, Harvard University, Cambridge, MA 02138; and [‡]Department of Biology, Kyushu University, Fukuoka 812-8581, Japan

Communicated by Bertram Kostant, Massachusetts Institute of Technology, Cambridge, MA, August 17, 2006 (received for review June 10, 2006)

Chronic myeloid leukemia (CML) is associated with the Philadelphia chromosome, which arises by a reciprocal translocation between chromosomes 9 and 22 and harbors the BCR-ABL fusion oncogene. It is unknown whether any other mutations are needed for the chronic phase of the disease. The CML incidence increases as a function of age with an exponent of ≈ 3 . A slope of 3 could indicate that there are two mutations, in addition to the Philadelphia translocation, that have not yet been discovered. In this work, we explore an alternative hypothesis: We study a model of cancer initiation requiring only a single mutation. A mutated cell has a net reproductive advantage over normal cells and, therefore, might give rise to clonal expansion. The cancer is detected with a probability that is proportional to the size of the mutated cell clone. This model has three waiting times: (i) the time until a mutated cell is produced, (ii) the time of clonal expansion, and (iii) the time until the clone is detected. Surprisingly, this simple process can give rise to cancer incidence curves with exponents up to 3. Therefore, the CML incidence data are consistent with the hypothesis that the Philadelphia translocation alone is sufficient to cause chronic phase CML.

cancer incidence | mathematical biology | stochastic dynamics

Chronic myeloid leukemia (CML) is a malignant clonal disorder of the hematopoietic system that leads to increased numbers of myelocytes, erythrocytes, and thrombocytes in peripheral blood (1, 2). The molecular hallmark of CML is the Philadelphia chromosome, first described as shortened chromosome 22 in 1960 (3) and then as a reciprocal t(9;22) translocation in 1973 (4). This abnormal chromosome is found in cells from the myeloid, erythroid, megakaryocytic, and B lymphoid lineages, indicating the presence of a “cancer stem cell” that is capable of producing several types of differentiated cancer cells (2). The Philadelphia chromosome is present in 95% of patients. The remaining 5% have complex or variant translocations that have the same result: fusion of the breakpoint cluster region (BCR) gene on chromosome 22 to the Ablson leukemia virus (ABL) gene on chromosome 9. The chimeric oncogene BCR-ABL encodes a constitutively active cytoplasmic tyrosine kinase. This protein activates growth and differentiation pathways in hematopoietic cells; its effectors include RAS, RAF, MYC, JUN, STAT, and phosphatidylinositol 3-kinase (1). BCR-ABL initiates a process to transform hematopoietic cells such that their growth and survival become independent of cytokines (5).

The question of whether BCR-ABL is necessary and sufficient to cause CML has been addressed in different ways. Several laboratories could reproduce a CML-like disease in mice expressing the BCR-ABL oncogene alone (6–8) or in combination with v-abl (9). Exposure to ionizing irradiation was shown to increase the risk of acquiring CML, but disease usually develops only after a prolonged latent period (10); this finding could indicate that several mutations are necessary for CML, but it also could mean that the Philadelphia-positive cell clone has a slow rate of expansion. Finally, $\approx 30\%$ of healthy individuals express the BCR-ABL oncogene at very low levels (11, 12). This observation, however, does not prove that additional genetic changes are necessary for CML because it can

alternatively be explained by the expression of BCR-ABL in non-self-renewing differentiated cells rather than hematopoietic stem cells. In the latter case, the continuous production of healthy hematopoietic cells will eventually replace the Philadelphia-positive cell clone and the disease will be “washed out” of the system (13). Hence, the experimental evidence of how many mutations are necessary to cause CML is inconclusive.

The multistep theory of carcinogenesis was inspired by the observation that the cancer incidence increases as a higher order function of age. In the 1950s, Nordling, Armitage, and Doll (14, 15) pointed out that the cancer incidence data could be explained by an increasing somatic mutation rate with age, or by the requirement of several events to cause cancer (16). In 1957, a simple deterministic model of two mutations was shown to produce incidence curves with slope six if the intermediate mutants lead to clonal expansion (17). Later, Moolgavkar *et al.* (16, 18, 19) introduced modeling of incidence curves based on multistate branching processes including cell proliferation and death.

Based on these insights, many mathematical models have been developed that investigate the number of mutations needed to cause particular kinds of cancer (19–22). A model fitted to the age-specific incidence of colorectal cancer proposes that two rare events and one high-frequency event suffice to initiate clonal expansion of the mutated stem cell, and only one more mutation is needed to give rise to an adenomatous polyp (19). The age-specific incidence of CML was used to calibrate a multistage model and predicts that three stem cell mutations are necessary for the chronic phase of the disease (20). These models neglect the population size and structure of the cells that are prone to accumulating mutations. However, the population genetics of susceptible cells must be considered if meaningful conclusions shall be drawn. For example, depending on the effective population size as compared with the mutation rate, it can take two, one, or even zero rate-limiting hits to accumulate two genetic alterations (23–26). The number of rate-limiting hits also can be larger than the actual number of mutations because of time delays due to clonal expansion (17). Therefore, the number of mutations necessary to cause a particular type of cancer cannot simply be read off the age-specific incidence data, and mathematical analyses must take into account population size and structure.

In the following, we offer a population genetics analysis of the dynamics of cancer initiation and its epidemiological consequences. Our model is motivated by CML but does not exclusively apply to this disease; rather, it is a general approach to studying many different types of cancer. We use CML as a specific example to show that a single mutation model can generate cancer incidence curves with an exponent of up to 3.

Author contributions: F.M. designed research; F.M., Y.I., and M.A.N. performed research; F.M. analyzed data; and F.M., Y.I., and M.A.N. wrote the paper.

The authors declare no conflict of interest.

Abbreviation: ALV, Ablson leukemia virus; BCR, breakpoint cluster region; CML, chronic myeloid leukemia; SEER, Surveillance, Epidemiology, and End Results.

[†]To whom correspondence should be addressed. E-mail: franziska.michor@harvard.edu.

© 2006 by The National Academy of Sciences of the USA




Fig. 1. The Moran process. Initially, all N cells are wild type and have a relative growth rate of 1. At each time step, a cell is chosen for reproduction proportional to fitness, and its daughter cell replaces another randomly chosen cell. The total number of cells remains strictly constant. Wild-type cells give rise to mutated cells at rate u per cell division. Mutated cells have a relative growth rate r . Cells divide every τ days.

Results and Discussion

Assume that a population of N (hematopoietic stem) cells proliferate according to the Moran process (27, 28): Initially, all cells are wild type. Cells divide every τ days. At each time step, a cell is chosen for reproduction proportional to fitness. The newly produced daughter cell replaces another randomly chosen cell (Fig. 1). Hence, the population size remains strictly constant. A wild-type cell gives rise to a mutated cell with probability u per cell division. Back mutation is neglected. A mutated cell has relative fitness r : if $r = 1$, then the mutation is neutral and the cell has the same growth rate as a wild-type cell; if $r > 1$, the mutant is advantageous, and if $r < 1$, the mutant is disadvantageous as compared with wild-type cells. We assume that the probability to diagnose the disease is linearly proportional to the number of mutated cells in the population. If there are x_1 mutant cells, the rate of detection is qx_1 . We perform exact numerical simulations of this process and compare the results with analytic approximations.

In *Methods*, we show that the probability of detecting the disease before time t is given by

$$P(t) = \int_0^t \left[1 - \left(1 + \frac{e^{cz} - 1}{N(1 - 1/r)} \right)^{-qN/c} \right] e^{-b(t-z)} b dz. \quad [1]$$

Here, $b = Nu(1 - 1/r)/\tau$ and $c = (r - 1)/\tau$. The stochastic process is characterized by three waiting times: (i) the waiting time until production of the first successful mutated cell is given by $1/b$; (ii) the time for clonal expansion of its lineage is given by $\ln[N(1 - 1/r)]/c$, and (iii) the time until detection of the disease is given by $1/qN$. This stochastic process can give rise to incidence curves with exponents up to 3. Define t_0 as the age at which we are interested in the slope of the incidence; for example, $t_0 = 50$ years. The factor b is always much less than the inverse of t_0 because otherwise, almost everybody would be diagnosed with cancer. Patients are diagnosed with a probability proportional to their number of leukemic stem cells, and if this probability is small, then the average time to detection might be >100 years. If $qN \gg 1/t_0$ and $c \gg 1/t_0$, then the incidence increases with an exponent of 1. In this case, the waiting time is dominated by the time it takes to produce the first successful mutant, and the time needed for clonal expansion and




Fig. 2. Simulation and theory. The figure shows the results of the exact stochastic computer simulation (triangles) and Eq. 1 (lines). Parameter values are mutation rate $u = 10^{-9}$, population size $N = 1,000$, relative fitness of mutated cells $r = 1.1$, mean cell generation time $\tau = 1$ day, and probability of detection $q/N = \alpha = 0.1$ for an exponent of 1 (a); $u = 10^{-5}$, $N = 2,000$, $r = 1.001$, $\tau = 100$, and $\alpha = 2 \times 10^{-5}$ for an exponent of 2 (b); and $u = 10^{-6}$, $N = 2,000$, $r = 1.01$, $\tau = 40$, and $\alpha = 2 \times 10^{-5}$ for an exponent of 3 (c).

detection is negligible and, hence, does not contribute to the slope. If $qN \ll 1/t_0$ and $c \gg 1/t_0$, then the slope is $2 + \alpha$, where α is between 0 and 1. Finally, if $qN \ll 1/t_0$ and $c \ll 1/t_0$, then the slope can take a value of 3. This contribution is only important if the time for clonal expansion is sufficiently long, i.e., if the mutated cell clone has a small fitness advantage. Therefore, a simple one-mutation model can give rise to incidence curves with an exponent of up to 3. Fig. 2 shows the exact stochastic simulation and Eq. 1 shows exponents 1–3.

The age-specific incidence data for CML were obtained from the Surveillance, Epidemiology, and End Results (SEER) registry, which covers $\approx 10\%$ of the U.S. population (www.seer.cancer.gov). In total, 5,256 CML cases were observed between 1973 and 2002, and these cases were recorded in age classes of five years (Table 1, columns 1 and 2). To calculate the probability to get CML per year from this data, some adjustments are made. First, the cases re-

